

The Leading Power Corrections to the Structure Functions

Xiaofeng Guo¹ and Jianwei Qiu^{2,3}

¹*Department of Physics and Astronomy, University of Kentucky,
Lexington, Kentucky 40506, USA*

²*Department of Physics and Astronomy, Iowa State University
Ames, Iowa 50011, USA*

³*Physics Department, Brookhaven National Laboratory
Upton, New York 11973-5000, USA
(October 29, 1998)*

Abstract

We show that when Bjorken x_B is large, the leading order twist-four contributions to the structure functions can be expressed in terms of the derivatives of the normal twist-two parton distributions. Our analytical results are not only consistent with the parameterized power corrections extracted from the data, but also predict the flavor dependence of the power corrections. We also discuss the impact of our analytical results on the extraction and the flavor separation of the parton distributions at large x .

PACS numbers: 12.38.Bx, 13.60.Hb

I. INTRODUCTION

Perturbative Quantum Chromodynamics (QCD) has been very successful in interpreting and predicting scattering cross sections at large momentum transfer. For hadronic collisions, a number of sets of reliable parton distributions are available for perturbative calculations of potential experimental signals [1]. However, recent data on inclusive high P_T jets from Fermilab [2] and high Q^2 events at HERA [3] remind us that there are still large uncertainties in the parton distributions at large x . The precise knowledge of parton distributions at large x is very important in search for signals of new physics at very high energies, which might be reached only by the partons with large momentum fraction x at the colliders.

According to the QCD perturbation theory, parton distributions at large x and high Q^2 can be derived from the parton distributions at large x and lower Q^2 by solving the DGLAP evolution equations. However, data from most physical observables are only sensitive to the parton distributions in the intermediate or small x region, because of the phase space integration and the fact that the parton distributions decrease very fast as x increases. It is believed that structure functions measured in deeply inelastic scattering (DIS) at large Bjorken x_B and low Q^2 might be the best source to get the direct information on the parton distributions at large x . But, at large x_B and low Q^2 , the structure functions receive significant power corrections [4], and these power corrections confuse the determination of the parton distributions.

In general, there are two types of power corrections, known as the kinematic and the dynamical power corrections. The kinematic power corrections is due to the fact that the nucleon has a finite mass, m_N . Such power corrections have been systematically discussed in Ref. [5]. On the other hand, the dynamical power corrections are the results of the interactions between quarks and gluons [5–7]. The dynamical power corrections are in general proportional to the multiparton correlation functions or the matrix elements of high twist operators, which are different from the normal parton distributions. Because of the large number of unknown multiparton correlation functions, the actual size of the dynamical power corrections to the structure functions have not been well understood theoretically. In this paper, we derive analytically the leading dynamical power corrections to the structure functions, and provide the numerical estimates of the power corrections. Since we will not discuss the kinematic power corrections in this paper, without any confusion, the words “power corrections” in the rest of this paper indicate the “dynamical power corrections”.

In a spin-averaged lepton-hadron deeply inelastic scattering, $l(k) + h(p) \rightarrow l(k') + X$, the hadronic tensor $W^{\mu\nu}(x_B, Q^2)$ can be decomposed as [8]

$$W^{\mu\nu}(x_B, Q^2) = - \left(g^{\mu\nu} - \frac{q^\mu q^\nu}{q^2} \right) F_1(x_B, Q^2) + \frac{1}{p \cdot q} \left(p^\mu - \frac{p \cdot q}{q^2} q^\mu \right) \left(p^\nu - \frac{p \cdot q}{q^2} q^\nu \right) F_2(x_B, Q^2), \quad (1)$$

where $q = k - k'$ is the four-momentum of the virtual photon, $q^2 = -Q^2$, and $F_i(x_B, Q^2)$ with $i = 1, 2$ are the structure functions. At the large enough Q^2 , the structure functions can be factorized into a convolution of the short-distance coefficient functions and nonperturbative matrix elements of multiparton fields [8],

$$F_i(x_B, Q^2) = \sum_{j,m} C_i^{(j,m)}(x_B, x_1, x_2, \dots, Q^2/\mu^2) \otimes T^{(j,m)}(x_1, x_2, \dots, \mu^2) \left(\frac{1}{Q^2} \right)^m, \quad (2)$$

where x_i with $i = 1, \dots$ are parton momentum fractions, μ^2 is the factorization scale and is often chosen to be Q^2 in DIS, $C^{(j,m)}$ are the perturbatively calculable coefficient functions, and $T^{(j,m)}$ are the corresponding nonperturbative matrix elements. In Eq. (2), $\sum_m$ runs from 0 to infinity, and $\sum_j$ sums over all possible types of nonperturbative matrix elements at a given power of m . The term with $m = 0$ is often known as the twist-2 or leading power contributions, and the terms with $m \neq 0$ are the high twist contributions or power corrections. The factorization presented in Eq. (2) is useful only when the terms of high power corrections can be neglected. When the physical scale, Q^2 , is large enough, the leading power contributions are often sufficient, and the structure functions can be expressed in terms of a few nonperturbative matrix elements, $T^{(j,0)}(x, Q^2)$. These matrix elements $T^{(j,0)}(x, Q^2)$ with $j = q, \bar{q}$ and g are known as the normal quark, antiquark, and gluon distributions, and have been well-determined for a wide range of x and Q^2 [1].

Working in the moment space of the structure functions, $F_i(n, Q^2) = \int_0^1 dx x^n F_i(x, Q^2)$, De Rújula, et al. [5] demonstrated that when the moment n is large enough,

$$F_2(n, Q^2) \sim \sum_m \left[\frac{n \mu_0^2}{Q^2} \right]^m T^m(n, Q^2), \quad (3)$$

with μ_0^2 a non-perturbative mass scale, and concluded that there exist a region of $n \sim Q^2/\mu_0^2$ where the power corrections are as important as the leading contributions. This conclusion is equivalent to say that in the x_B -space, when $x_B \rightarrow 1$, high order terms in the expansion in Eq. (2) become important even if Q^2 is large. Physically, when $x_B \rightarrow 1$, the invariant mass of the photon-hadron system in DIS approaches to the resonance region, and the perturbative expansion in Eq. (2) is then invalid.

In order to avoid the resonances, we have to require the invariant mass of the photon-hadron system [9,10]

$$W^2 \equiv (p + q)^2 = m_N^2 + \frac{1 - x_B}{x_B} Q^2 > 2 \text{ GeV}^2. \quad (4)$$

On the other hand, in order to extract the information on parton distributions at large x_B and low Q^2 , we have to work in a region where W^2 is small and not too far away from the resonances, and will have to deal with the power corrections. The purpose of this paper is to get a good estimate of the size of the leading power corrections and to discuss its impact on the extraction of the reliable twist-2 parton distributions at large x .

By fitting DIS data in large x_B region, it has been found [4,11] that the $1/Q^2$ dynamical power corrections to the structure functions can be parameterized in terms of the normal parton distributions,

$$F_2(x_B, Q^2) \approx \left(1 + \frac{h(x_B)}{Q^2} \right) F_2(x_B, Q^2)_{\text{LT}}, \quad (5)$$

where $F_2(x_B, Q^2)_{\text{LT}}$ includes all leading twist (or leading power) contributions as well as the target mass corrections to the full structure function $F_2(x_B, Q^2)$. The $h(x)$ in Eq. (5) is a phenomenological fitting function, and is parameterized as [11]

$$h(x_B) = a \left(\frac{x_B^b}{1 - x_B} - c \right), \quad (6)$$

where a, b and c are constant fitting parameters. Because of the $1/(1 - x_B)$ dependence in Eq. (6), it is clear that the dynamical power corrections in Eq. (5) become more important when x_B increases. It has been argued that the simple parameterization in Eq. (6) can provide a very good fit to the existing DIS data on proton as well as deuteron targets (with slightly different fitting parameters), and demonstrated that the power corrections are important for extracting the correct parton distributions, such as d/u ratio, in large x region [4,11].

On the other hand, a complete expression of the $1/Q^2$ power corrections to the structure functions at the leading order of α_s were derived more than fifteen years ago [12,13]. As expected from Eq. (2), the $1/Q^2$ corrections are proportional to the matrix elements of four-parton operators, which are very different from the operators defining the normal parton distributions. The success of the phenomenological parameterization in Eq. (5) raises a question: under what conditions, one can derive the approximate expression in Eq. (5), or a similar expression, from the complete $1/Q^2$ power corrections at the leading order of α_s , which were derived in Ref. [12,13].

In this paper, we show that when x_B is large, the $1/Q^2$ power corrections to the structure functions at the leading order of α_s can be approximated as

$$F_2(x_B, Q^2) \approx \sum_{q, \bar{q}} e_q^2 x_B \left\{ 1 + \frac{1}{Q^2} \left[\langle D_T^2 \rangle \left(-x_B \frac{d}{dx_B} \right) + 4 \langle m_T^2 \rangle \right] \right\} q(x_B, Q^2), \quad (7)$$

where $m_N^2 = p^2$ is the nucleon mass, $\langle D_T^2 \rangle$ and $\langle m_T^2 \rangle$ are the averaged values of D_T^2 and m_T^2 , which will be defined in the next section. Numerically, we demonstrate that our analytical expressions for the leading power corrections in Eq. (7) have the same x_B -dependence as the phenomenological parameterization given in Eqs. (5) and (6), and therefore, we expect Eq. (7) to fit the data as well. Instead of three unknown parameters, a, b and c in Eq. (6), our analytical result in Eq. (7) have only two parameters, and both of them have clear physical interpretations. Furthermore, because of the scaling violation of the quark distributions, the derivative in Eq. (7) gives the natural explanation for the flavor dependence (or target dependence) of the fitting parameters in Eq. (6) [11]. In addition, the derivative provides some extra logarithmic Q^2 -dependence for the power corrections.

Our result in Eq. (7) demonstrates that although the power corrections to the structure functions in DIS are in general very complicated, the leading contributions in α_s at large x_B are approximately given by the derivatives of the normal quark distributions. Such a simple analytical expression will be very useful for the QCD global analysis of extracting reliable leading twist parton distributions.

The rest of our paper is organized as follows. For the completeness, we briefly derive the leading twist-4 contributions to the structure functions in Sec. II. In Sec. III, we identify the necessary approximations, and derive the main result of this paper, Eq. (7), from the complete leading power corrections obtained in Sec. II. Finally, in Sec. IV, we numerically compare our analytical expression in Eq. (7) with the phenomenological parameterization in Eqs. (5) and (6), and estimate the values of the parameters $\langle D_T^2 \rangle$ and $\langle m_T^2 \rangle$ in Eq. (7). We also discuss the possible impact of our analytical results of the leading power corrections to the extraction and the flavor separation of parton distributions at large x .

II. POWER CORRECTIONS IN DIS AT LEADING ORDER IN α_s

The complete $1/Q^2$ power corrections to the structure functions at the leading order of α_s can be separated into two groups: one is proportional to the two-quark-two-gluon matrix elements (or correlation functions), and the other is proportional to the four-quark matrix elements [12,13]. Because of Fermi statistics and the fact that sea quark distributions are smaller than the gluon distributions, contributions from the four-quark subprocesses in large x_B region should be smaller than that from the quark-gluon subprocesses. In this section, we neglect the four-quark contributions, and drive the leading order $1/Q^2$ contributions to the DIS structure functions from the quark-gluon subprocess, from which the main result of this paper, Eq. (7), will be deduced in the next section.

For the definiteness of our derivations, we choose the following photon-hadron frame

$$\begin{aligned} q^\mu &= -x_B p \cdot n \bar{n}^\mu + \frac{Q^2}{x_B p \cdot n} n^\mu , \\ p^\mu &= p \cdot n \bar{n}^\mu , \end{aligned} \quad (8)$$

where we have neglected the hadron mass for the dynamical power corrections. In Eq. (8), the four-vector, $\bar{n}^\mu$ and n^μ are defined as

$$\begin{aligned} \bar{n}^\mu &\equiv (\bar{n}^+, \bar{n}^-, \bar{n}_T) = (1, 0, 0_T) , \\ n^\mu &= (0, 1, 0_T) , \\ \bar{n}^2 &= 0, \quad n^2 = 0, \quad \text{and} \quad \bar{n} \cdot n = 1 . \end{aligned} \quad (9)$$

In terms of the $\bar{n}^\mu$ and n^μ , we can reexpress the hadronic tensor in Eq. (1) as

$$W^{\mu\nu}(x_B, Q^2) = e_L^{\mu\nu} F_L(x_B, Q^2) + e_T^{\mu\nu} F_1(x_B, Q^2) , \quad (10)$$

where the longitudinal and transverse tensors are defined as

$$\begin{aligned} e_L^{\mu\nu} &\equiv \frac{1}{2Q^2} \left(x_B p \cdot n \bar{n}^\mu + \frac{Q^2}{2x_B p \cdot n} n^\mu \right) \left(x_B p \cdot n \bar{n}^\nu + \frac{Q^2}{2x_B p \cdot n} n^\nu \right) , \\ e_T^{\mu\nu} &\equiv \bar{n}^\mu n^\nu + n^\mu \bar{n}^\nu - g^{\mu\nu} \equiv d^{\mu\nu} , \end{aligned} \quad (11)$$

with $e_{L\mu\nu} e_L^{\mu\nu} = 1/4$, $e_{T\mu\nu} e_T^{\mu\nu} = 2$, and $e_{L\mu\nu} e_T^{\mu\nu} = 0$. The longitudinal structure function, $F_L(x_B, Q^2)$ in Eq. (10), is defined as usual, $F_L = F_2/x_B - 2F_1$. From Eq. (10), the DIS structure functions can be extracted from the hadronic tensor as

$$\begin{aligned} F_1(x_B, Q^2) &= \frac{1}{2} e_T^{\mu\nu} W_{\mu\nu}(x_B, Q^2) \\ F_L(x_B, Q^2) &= 4 e_L^{\mu\nu} W_{\mu\nu}(x_B, Q^2) . \end{aligned} \quad (12)$$

As derived in Ref. [13], the $1/Q^2$ power corrections to the structure functions from the quark-gluon subprocesses can be factorized, as shown in Fig. 1,

$$W^{\mu\nu}(x_B, Q^2) \Big|_{1/Q^2} = \frac{1}{Q^2} \int dx dx_1 dx_2 \text{Tr} \left[C_{\alpha\beta}^{\mu\nu}(x_B, x, x_1, x_2) T^{\alpha\beta}(x, x_1, x_2, Q^2) \right] , \quad (13)$$

where "Tr" is the spinor trace for fermions. The partonic part, $C_{\alpha\beta}^{\mu\nu}(x_B, x, x_1, x_2)$ in above equation, is given by the top parts of the Feynman diagrams in Fig. 1. The blobs in Fig. 1a are given by the diagrams in Fig. 2a, and the blobs in Fig. 1b and Fig. 1c are given by the diagrams in Fig. 2b. In Fig. 2, the fermion lines with a short bar indicates that the propagator includes only the short-distance contact term, as defined in Ref. [13]. The quark-gluon correlation functions, $T^{\alpha\beta}(x, x_1, x_2, Q^2)$ in Eq. (13), is expressed in terms of the four-parton matrix element,

$$T^{\alpha\beta}(x, x_1, x_2, Q^2) = \int \frac{p^+ dy^-}{2\pi} \frac{p^+ dy_1^-}{2\pi} \frac{p^+ dy_2^-}{2\pi} e^{ix_1 p^+ y^-} e^{i(x-x_1)p^+ y_1^-} e^{-i(x-x_2)p^+ y_2^-} \\ \times \langle p | \bar{\psi}(0) D_T^\beta(y_2^-) D_T^\alpha(y_1^-) \psi(y^-) | p \rangle, \quad (14)$$

where $D^\alpha(y) \equiv \partial^\alpha + ig T^B A_B^\alpha(y)$ is the covariant derivative, which differs by a factor of i from the definition used in Refs. [12,13]. In Eq. (14), $D_T^\alpha \equiv (g^{\alpha\beta} - \bar{n}^\alpha n^\beta) D_\beta$, and the line-integrals between the fields are omitted [13].

To complete the factorization between the partonic parts and the twist-4 matrix elements, we need to separate the spinor trace, as well as the Lorentz indices linking between the $C_{\alpha\beta}^{\mu\nu}$ and $T^{\alpha\beta}$ in Eq. (13). Such factorization can be achieved by decomposing the $T^{\alpha\beta}$ as

$$T^{\alpha\beta} = \frac{1}{2} \gamma \cdot p d^{\alpha\beta} T + \frac{1}{2} \gamma \cdot p \gamma_5 (i\epsilon^{\alpha\beta}) \tilde{T} + \dots, \quad (15)$$

where T and $\tilde{T}$ are scalar functions of x, x_1, x_2 and Q^2 , and "... represents terms that are further power suppressed. In Eq. (15), the symmetric tensor $d^{\alpha\beta}$ is defined in Eq. (11) and the antisymmetric tensor $\epsilon^{\alpha\beta} = \epsilon^{\alpha\beta\mu\nu} \bar{n}_\mu n_\nu$. From Eqs. (12) and (13), we can derive the $1/Q^2$ contributions to the structure functions at the leading order in α_s by calculating the Feynman diagrams in Fig. 2. Substituting Eq. (15) into Eq. (13), and contracting $e_T^{\mu\nu}$ with the hadronic tensor $W_{\mu\nu}(x_B, Q^2)|_{1/Q^2}$, we obtain in the light-cone gauge ($n \cdot A = 0$),

$$e_T^{\mu\nu} W_{\mu\nu}^{(a)}(x_B, Q^2) \Big|_{1/Q^2} = 0, \quad (16a)$$

$$e_T^{\mu\nu} W_{\mu\nu}^{(b)}(x_B, Q^2) \Big|_{1/Q^2} = -e_q^2 \frac{2x_B}{Q^2} \int dx_1 dx_2 \frac{\delta(x_1 - x_B)}{x_2 - x_B} T_q(x_2, x_1, Q^2), \quad (16b)$$

$$e_T^{\mu\nu} W_{\mu\nu}^{(c)}(x_B, Q^2) \Big|_{1/Q^2} = -e_q^2 \frac{2x_B}{Q^2} \int dx_1 dx_2 \frac{\delta(x_2 - x_B)}{x_1 - x_B} T_q(x_2, x_1, Q^2), \quad (16c)$$

where e_q is the quark(or antiquark)'s fractional charge of flavor q , and the superscripts (a), (b) and (c) correspond to the contributions from diagrams Figs. 2a, 2b and 2c, respectively. The quark-gluon correlation function $T_q(x_2, x_1, Q^2)$ is defined as

$$T_q(x_2, x_1, Q^2) = \frac{1}{4} \int \frac{dy^-}{2\pi} \frac{p^+ dy_1^-}{2\pi} e^{ix_1 p^+ y^-} e^{i(x_2 - x_1)p^+ y_1^-} \\ \times \langle p | \bar{\psi}_q(0) \gamma^+ (d_{\alpha\beta} D^\alpha(y_1^-) D^\beta(y_1^-)) \psi_q(y^-) | p \rangle. \quad (17)$$

Substituting Eq. (16) into Eq. (12), we obtain

$$F_1(x_B, Q^2) \Big|_{1/Q^2} = \sum_q e_q^2 \frac{x_B}{Q^2} \int dx_1 dx_2 \left[\frac{\delta(x_2 - x_B) - \delta(x_1 - x_B)}{x_2 - x_1} \right] T_q(x_2, x_1, Q^2). \quad (18)$$

Similarly, by contracting $e_L^{\mu\nu}$ to the hadronic tensor $W_{\mu\nu}(x_B, Q^2)$, we obtain

$$e_L^{\mu\nu} W_{\mu\nu}^{(a)}(x_B, Q^2) \Big|_{1/Q^2} = e_q^2 \frac{2}{Q^2} \int dx \delta(x - x_B) T_q(x, Q^2) , \quad (19a)$$

$$e_L^{\mu\nu} W_{\mu\nu}^{(b)}(x_B, Q^2) \Big|_{1/Q^2} = 0 , \quad (19b)$$

$$e_L^{\mu\nu} W_{\mu\nu}^{(c)}(x_B, Q^2) \Big|_{1/Q^2} = 0 , \quad (19c)$$

where the quark-gluon correlation function $T_q(x, Q^2)$ is defined as

$$T_q(x, Q^2) = -\frac{1}{4} \int \frac{dy^-}{2\pi} e^{ixp^+ y^-} \langle p | \bar{\psi}_q(0) \gamma_\alpha D_T^\alpha(0) \gamma^+ \gamma_\beta D_T^\beta(y^-) \psi_q(y^-) | p \rangle . \quad (20)$$

Combining Eqs. (12) and (19), we have

$$F_L(x_B, Q^2) \Big|_{1/Q^2} = \sum_q e_q^2 \frac{8}{Q^2} T_q(x_B, Q^2) . \quad (21)$$

From Eqs. (18) and (21), we obtain the leading order $1/Q^2$ contributions from quark-gluon subprocess to the structure function $F_2(x_B, Q^2)$ as

$$\begin{aligned} F_2(x_B, Q^2) \Big|_{1/Q^2} = \frac{1}{Q^2} \sum_q e_q^2 x_B \Big(2x_B \int dx_1 dx_2 \left[\frac{\delta(x_2 - x_B) - \delta(x_1 - x_B)}{x_2 - x_1} \right] T_q(x_2, x_1, Q^2) \\ + 8 \int dx \delta(x - x_B) T_q(x, Q^2) \Big) . \end{aligned} \quad (22)$$

Our results given in Eqs. (18), (21) and (22) are consistent with those in Refs. [12,13], after taking into account the difference in the definition of the covariant derivative (a factor of “ i ”) and an extra “ -1 ” in the definition of $T_q(x, Q^2)$ in Eq. (20).

III. LEADING POWER CORRECTIONS AT LARGE x_B

In this section, we identify the approximations that enable us to derive Eq. (7) from Eq. (22), which represents the complete $1/Q^2$ contributions to $F_2(x_B, Q^2)$ from the leading order quark-gluon subprocesses.

From the phenomenological fitting of the dynamical power corrections [11], $1/Q^2$ contributions to the structure functions are most important in large x_B region, because of the $1/(1-x_B)$ factor in Eq. (6). From the complete leading order $1/Q^2$ contributions in Eq. (22), it is nontrivial to conclude that the $1/(1-x_B)$ behavior exists. However, under some simple approximations, we demonstrate below that Eq. (22) has indeed the $1/(1-x_B)$ behavior when x_B is large.

The second term in Eq. (22) is simple and directly proportional to the quark-gluon correlation function $T_q(x_B, Q^2)$ in Eq. (20). To see if the $T_q(x_B, Q^2)$ has the $1/(1-x_B)$ behavior, we simplify the operator expression of the $T_q(x_B, Q^2)$ by using the equation of motion, $\gamma_\alpha D^\alpha(y) \psi_q(y) = 0$, and we have

$$\begin{aligned}
T_q(x_B, Q^2) &= -\frac{1}{4} \int \frac{dy^-}{2\pi} e^{ix_B p^+ y^-} \langle p | \bar{\psi}_q(0) \gamma^- D^+(0) \gamma^+ \gamma^- D^+(y^-) \psi_q(y^-) | p \rangle \\
&= \frac{(x_B)^2}{2} \int \frac{p^+ dy^-}{2\pi} e^{ix_B p^+ y^-} \langle p | \bar{\psi}_q(0) [p^+ \gamma^-] \psi_q(y^-) | p \rangle
\end{aligned} \tag{23}$$

where the light-cone gauge $n \cdot A = 0$ and $D_T^\alpha = (g^{\alpha\beta} - \bar{n}^\alpha n^\beta) D_\beta$ were used. On the other hand, the normal twist-2 quark distribution can be expressed in the light-cone gauge as

$$\begin{aligned}
q(x_B, Q^2) &= \frac{1}{2} \int \frac{dy^-}{2\pi} e^{ix_B p^+ y^-} \langle p | \bar{\psi}_q(0) \gamma^+ \psi_q(y^-) | p \rangle \\
&= \frac{1}{m_N^2} \int \frac{p^+ dy^-}{2\pi} e^{ix_B p^+ y^-} \langle p | \bar{\psi}_q(0) [p^- \gamma^+] \psi_q(y^-) | p \rangle
\end{aligned} \tag{24}$$

where $m_N^2 \equiv 2p^+ p^-$ is the nucleon mass square. In order to relate the correlation function $T_q(x_B, Q^2)$ to the quark distribution $q(x_B, Q^2)$, we estimate the difference between the matrix elements $\langle p | \bar{\psi}_q(0) [p^+ \gamma^-] \psi_q(y^-) | p \rangle$ in Eq. (23) and $\langle p | \bar{\psi}_q(0) [p^- \gamma^+] \psi_q(y^-) | p \rangle$ in Eq. (24) as follows. For free field operator ψ_q 's on free light quark states, the difference between these two matrix elements is the difference between $[p^+ k^-]$ and $[p^- k^+]$ with k the light quark momentum. In our frame, we can choose

$$p^- = m_N^2/(2p^+) \quad \text{and} \quad k^- = m_T^2/(2k^+) \quad \text{and} \quad k^+ = x_B p^+, \tag{25}$$

with the transverse mass of the light quark to be $m_T^2 \equiv m_q^2 + k_T^2$. If we *assume* the free field relation can be generalized to the full fields, we obtain

$$\begin{aligned}
&\int \frac{p^+ dy^-}{2\pi} e^{ix_B p^+ y^-} \langle p | \bar{\psi}_q(0) [p^+ \gamma^-] \psi_q(y^-) | p \rangle \\
&\approx \frac{\langle m_T^2 \rangle}{m_N^2 (x_B)^2} \int \frac{p^+ dy^-}{2\pi} e^{ix_B p^+ y^-} \langle p | \bar{\psi}_q(0) [p^- \gamma^+] \psi_q(y^-) | p \rangle.
\end{aligned} \tag{26}$$

Substituting Eq. (26) into Eqs. (23) and (24), we derive

$$T_q(x_B, Q^2) \approx \frac{1}{2} \langle m_T^2 \rangle q(x_B, Q^2). \tag{27}$$

We emphasize here that the importance of Eq. (27) is not how accurate the approximation is, rather it indicates that the second term in Eq. (22) does not have the $1/(1-x_B)$ enhancement.

The first term in Eq. (22) depends on the correlation function $T_q(x_2, x_1, Q^2)$, which can be sketched in Fig. 3, and be reexpressed as

$$\begin{aligned}
T_q(x_2, x_1, Q^2) &= \frac{1}{2} \int \frac{dy^-}{2\pi} e^{ix_1 p^+ y^-} \langle p | \bar{\psi}_q(0) \gamma^+ \\
&\quad \times \left[\frac{1}{2} \int \frac{p^+ dy_1^-}{2\pi} e^{i(x_2 - x_1) p^+ y_1^-} d_{\alpha\beta} D^\alpha(y_1^-) D^\beta(y_1^-) \right] \psi_q(y^-) | p \rangle.
\end{aligned} \tag{28}$$

Because of the factor $[\delta(x_2 - x_B) - \delta(x_1 - x_B)]/(x_2 - x_1)$ in Eq. (22), $x_1 \sim x_2 \sim x_B$ in $T_q(x_2, x_1, Q^2)$ when x_B is large, and the effective ‘‘gluons’’ in Fig. 3 are soft. We then assume that the effective ‘‘gluon’’ operator $d_{\alpha\beta} D^\alpha(y_1^-) D^\beta(y_1^-)$ in Eq. (28) is not sensitive to

the position variable y_1^- , and further approximate this operator by an averaged expectation value as

$$d_{\alpha\beta} D^\alpha(y_1^-) D^\beta(y_1^-) \approx \langle d_{\alpha\beta} D^\alpha(0) D^\beta(0) \rangle \equiv \langle D_T^2 \rangle . \quad (29)$$

More discussion on this assumption will be given in the next section. With the approximation in Eq. (29), we can rewrite the correlation function $T_q(x_2, x_1, Q^2)$ as

$$\begin{aligned} T_q(x_2, x_1, Q^2) &\approx \frac{1}{2} \int \frac{dy^-}{2\pi} e^{ix_1 p^+ y^-} \langle p | \bar{\psi}_q(0) \gamma^+ \left[\frac{1}{2} \langle D_T^2 \rangle \delta(x_2 - x_1) \right] \psi_q(y^-) | p \rangle , \\ &= \left[\frac{1}{2} \langle D_T^2 \rangle \delta(x_2 - x_1) \right] q(x_1, Q^2) , \end{aligned} \quad (30)$$

where $q(x_1, Q^2)$ is the twist-2 quark distribution defined in Eq. (24). Using Eq. (30), we derive

$$\begin{aligned} &\int dx_1 dx_2 \left[\frac{\delta(x_2 - x_B) - \delta(x_1 - x_B)}{x_2 - x_1} \right] T_q(x_2, x_1, Q^2) \\ &\approx \frac{1}{2} \langle D_T^2 \rangle \int_0^1 dx_1 \delta'(x_1 - x_B) q(x_1, Q^2) \\ &= \frac{1}{2} \langle D_T^2 \rangle \left(-\frac{d}{dx_B} \right) q(x_B, Q^2) . \end{aligned} \quad (31)$$

Substituting Eqs. (27) and (31) into Eq. (22), we obtain the leading power corrections to the structure function $F_2(x_B, Q^2)$ as

$$F_2(x_B, Q^2) \Big|_{1/Q^2} \approx \frac{1}{Q^2} \sum_q e_q^2 x_B \left[\langle D_T^2 \rangle \left(-x_B \frac{d}{dx_B} \right) + 4 \langle m_T^2 \rangle \right] q(x_B, Q^2) . \quad (32)$$

Combining this leading power corrections and the leading order twist-2 contributions to $F_2(x_B, Q^2)$, we obtain our main result, Eq. (7).

Although Eq. (32) looks very different from the phenomenological parameterization in Eq. (6), we show below that Eqs. (32) and (6) have the same analytical behavior as $x_B \rightarrow 1$. When x_B is large, the quark distribution have the following behavior

$$q(x_B, Q^2) \Rightarrow (1 - x_B)^{n_q(Q^2)} \quad \text{as } x_B \rightarrow 1 , \quad (33)$$

with a positive and real parameter n_q . In general, $n_q(Q^2)$ has the logarithmic Q^2 -dependence due to the scaling violation of the parton distributions. For the valence quark distributions, $n_q \sim 3$. Consequently, we have

$$-x_B \frac{d}{dx_B} q(x_B) = n_q(Q^2) \frac{x_B}{1 - x_B} q(x_B, Q^2) + \text{terms without } \frac{1}{1 - x_B} . \quad (34)$$

Substituting Eq. (34) into Eq. (32), naturally, we derive the leading $1/(1 - x_B)$ behavior,

$$F_2(x_B, Q^2) \Big|_{1/Q^2} \approx \sum_q e_q^2 x_B q(x_B, Q^2) \left[\frac{\langle D_T^2 \rangle}{Q^2} n_q(Q^2) \frac{x_B}{1 - x_B} + \text{terms without } \frac{1}{1 - x_B} \right] , \quad (35)$$

which is consistent with the phenomenological parameterization in Eq. (6) that was introduced in Refs. [4,11] after fitting the existing DIS data.

IV. NUMERICAL RESULTS AND CONCLUSIONS

In this section, we demonstrate numerically that as a function of x_B , our analytical expression in Eq. (7) and the phenomenological parameterization defined in Eqs. (5) and (6) have almost identical behavior, even though they look very different.

In the following numerical estimates, we use CTEQ3L parton distributions of Ref. [14] for the quark distributions in the proton. In addition, in order to avoid the resonance region and to test the perturbative power expansion of the structure functions, we keep $W^2 \geq 2 \text{ GeV}^2$ in our calculations.

In Fig. 4, we plot the structure function $F_2(x_B, Q^2)$ as a function of x_B for $Q^2 = 4$ and 9 GeV^2 . The solid line is the lowest order contribution, $F_2^{\text{LO}}(x_B, Q^2) = \sum_q e_q^2 x_B q(x_B, Q^2)$. The dotted and dashed lines are given by the expressions in Eqs. (5) and (7), respectively, with $F_2^{\text{LO}}(x_B, Q^2)$ for the $F_2(x_B, Q^2)_{\text{LT}}$ in Eq. (5). For the fitting parameters, a, b and c in the phenomenological parameterization $h(x)$, we use the following values $a = 0.50$, $b = 3.2$ and $c = 0.11$, which are given in Ref. [11]. For our analytical expression in Eq. (7), there are two unknown parameters $\langle D_T^2 \rangle$ and $\langle m_T^2 \rangle$. We choose $\langle D_T^2 \rangle = 0.1 \text{ GeV}^2$ to be consistent with the overall size of the power corrections. The transverse mass square $\langle m_T^2 \rangle$ for the light quarks should be of the order of the intrinsic transverse momentum square, and therefore, $\langle m_T^2 \rangle \geq \Lambda_{\text{QCD}}^2$. The actual size of $\langle m_T^2 \rangle$ should depend on the collision energy W^2 due to the soft gluon shower from the incoming quark line [15]. Within our approximation, different choices for the value of $\langle m_T^2 \rangle$ represents a constant shift of the size of the leading power corrections in $(F_2 - F_2^{\text{LO}})/F_2^{\text{LO}}$. In the rest of our discussions, we choose $\langle m_T^2 \rangle$ to be zero for a close exam of the derivative term. In this paper, we are more interested in understanding the functional dependence of the power corrections than trying to extract the best values of the $\langle D_T^2 \rangle$ and $\langle m_T^2 \rangle$ by fitting any data. We defer the detailed global QCD analysis for extracting the parton distributions from the data to another publication.

Because of the steep falling parton distributions when x_B increases, we plot the relative power corrections to the structure function, $(F_2 - F_2^{\text{LO}})/F_2^{\text{LO}}$, as a function of x_B in Fig. 5. As shown in Fig. 5, the leading order power corrections are very significant and can be as large as 30% in large x_B region. In addition, with only one parameter $\langle D_T^2 \rangle$ (because $\langle m_T^2 \rangle$ is set to be zero here), our analytical result in Eq. (7) is remarkably consistent with the parameterization extracted from the data. The difference between the solid and the dashed curves in Fig. 5 cannot be separated by the existing data. Such consistency on x_B -dependence is already a nontrivial test for the QCD perturbation theory.

As shown in last section, the leading power corrections at large x_B are given by the derivatives of the normal quark distributions. It is the derivative that makes the power corrections different for quark distributions of different flavors. We introduce

$$\Delta q(x_B, Q^2) \equiv \frac{1}{Q^2} \left[\langle D_T^2 \rangle \left(-x_B \frac{d}{dx_B} \right) + 4 \langle m_T^2 \rangle \right] q(x_B, Q^2), \quad (36)$$

for quark flavor q . In Figs. 6 and 7, we plot the relative power corrections $\Delta q/q$ as a function of x_B for the “up” and “down” quark, respectively. Because the “down” quark distribution falls off faster than the “up” quark distribution in the CTEQ3L parton distributions, the “down” quark obtains a larger power correction than what the “up” quark gets. Such difference is clearly evident in Figs. 6 and 7, and is very significant when x_B is large. The

power corrections derived here also receive some logarithmic Q^2 -dependence through the derivative of the parton distributions, due to the scaling violation of the twist-2 parton distributions.

As demonstrated above, the power corrections to the structure functions are not only significant at large x_B , but also sensitive to the quark flavors. It is such flavor difference of the power corrections that makes the extraction and the flavor separation of the parton distributions nontrivial, in particular, in large x_B region. Recently, a lot of attention have been devoted to the ratio of d/u [4]. We defer the detailed global QCD analysis for the extraction and the flavor separation of the parton distributions to another publication.

In summary, we derive the analytical expressions for the leading power corrections to the DIS structure function, given in Eq. (32). We made two main assumptions in our derivation, which are introduced in Eqs. (26) and (29), respectively. Since the $1/(1-x_B)$ enhancement of the power corrections is from the term proportional to the quark-gluon correlation $T_q(x_2, x_1, Q^2)$, the assumption in Eq. (29) is more crucial than the other. In general, it may be possible that the $\langle D_T^2 \rangle$ is not a constant independent of the position (or equivalently the value of x_B). However, as long as any part of the $\langle D_T^2 \rangle$ is not proportional to $(1-x_B)$, we will have the $1/(1-x_B)$ enhancement. We believe that at large x_B , at least the “soft” gluon part of the averaged value of the covariant derivative square should not be sensitive the location inside the nucleon, even though the averaged value of the transverse momentum square may be proportional to $(1-x_B)$. Therefore, we conclude that the leading order power corrections to the structure function $F_2(x_B, Q^2)$ should have the $1/(1-x_B)$ enhancement. and our analytical result, Eq. (7) with two fitting parameters $\langle D_T^2 \rangle$ and $\langle m_T^2 \rangle$, should predicts the correct *leading* x_B -dependence of the power corrections at large x_B . Because it has only two unknown parameters, our result has a good predicting power and can be tested at CEBAF as well as future fixed target programs at Fermilab. It is also useful for the QCD global analysis for extracting parton distributions in large x_B region. In addition, our approach can be applied to the Drell-Yan process to study the leading power corrections there [16].


ACKNOWLEDGMENT

We are pleased to acknowledge useful discussions with C. Keppel and W. Zhu. This work was supported in part by the U.S. Department of Energy under Grant Nos. DE-FG02-87ER40731 and DE-FG02-96ER40989, and under the Contract No. DE-AC02-98CH10886.


REFERENCES

- [1] For a review and detailed references, see J.F. Owens and W.K. Tung, *Ann. Rev. Nucl. Sci.* **42**, 291 (1992).
- [2] F. Abe *et al.*, CDF Collaboration, *Phys. Rev. Lett.* **77**, 438 (1996); G. Blazey *et al.*, D0 Collaboration,
- [3] C. Adloff *et al.*, *Z. Phys.* **C74**, 191 (1997); J. Breitweg *et al.*, *Z. Phys.* **C74**, 207 (1997).
- [4] For example, U.K. Yang and A. Bodek, hep-ph/9809480, and references there in.
- [5] A. De Rújula, H. Georgi, and H.D. Politzer, *Ann. Phys. (N.Y.)* **103**, 315 (1977).
- [6] R.L. Jaffe and M. Soldate, *Phys. Lett.* **B105**, 467 (1981); *Phys. Rev.* **D26**, 49 (1982).
- [7] M. Dasgupta and B.R. Webber, *Phys. Lett.* **B382**, 273 (1996).
- [8] G. Sterman *et al.*, *Rev. Mod. Phys.* **67**, 157 (1995).
- [9] A. Bodek *et al.*, *Phys. Rev.* **D20**, 1471 (1979).
- [10] X. Ji and P. Unrau, *Phys. Rev.* **D52**, 72 (1996).
- [11] U.K. Yang and A. Bodek, hep-ph/9806458.
- [12] R.K. Ellis, W. Furmanski, and R. Petronzio, *Nucl. Phys.* **B212**, 29 (1983).
- [13] J.-W. Qiu, *Phys. Rev.* **D42**, 30, (1990).
- [14] H. L. Lai, *et al.*, *Phys. Rev.* **D51**, 4763 (1995); and references therein.
- [15] J. Huston *et al.*, *Phys. Rev.* **D51**, 6139 (1995).
- [16] J.-W. Qiu and G. Sterman, *Nucl. Phys.* **B353**, 105 (1991).


FIGURES



(a)



(b)



(c)

Fig. 1

FIG. 1. Factorized diagrams for the leading order power corrections from the quark-gluon subprocesses.




Fig. 2

FIG. 2. Feynman diagrams for the blobs in Fig. 1. The wave lines and curly lines represent the virtual photons and gluons, respectively. The solid lines are for the quarks, and the line with a short bar represents the contact term of the propagator [13].






Fig. 3

FIG. 3. Diagram represents the quark-gluon correlation function $T_q(x_2, x_1, Q^2)$.



(a)


Fig. 4



(b)


Fig. 4

FIG. 4. Structure function $F_2(x_B, Q^2)$ as a function of x_B at 4 GeV² (a), and 9 GeV² (b).



(a)

Fig. 5



(b)

Fig. 5

FIG. 5. Relative power corrections to the structure function $F_2(x_B, Q^2)$ as a function of x_B at 4 GeV² (a), and 9 GeV² (b).




Fig. 6

FIG. 6. Relative power corrections to the “up” quark distribution at $Q^2 = 4 \text{ GeV}^2$ and $Q^2 = 9 \text{ GeV}^2$, respectively.




Fig. 7

FIG. 7. Relative power corrections to the “down” quark distribution at $Q^2 = 4 \text{ GeV}^2$ and $Q^2 = 9 \text{ GeV}^2$, respectively.